

Mixing Instructions

With **KBC 35/30 SA & KLC 10**
 Country **Uganda**
 Specie **Layers**
 Batch weight **1,000 kg**

9201-9211 Economy



Version: 5/31/2023

Code	Ingredient	Unit (kg)	960710 Pre-Starter 0-7 days	201-9201 Starter 1-6 weeks	202-9201 Grower 7-12 weeks	204-9211 Developer 13-17 weeks	205-9211 Pre layer 18-19 weeks (till 5% production)	206-9211 Pre Peak 20-28 weeks	207-9211 Phase 1 29-55 weeks	208-9211 Phase 2 56 weeks - end
960710	Galdus Layer	Min. 100gr/DOC								
KN627C	MAIZE			550	530	550	450	470	460	440
969452	KBC 35/30 SA			350	300					
KN673C	WHEAT BRAN / POLLARD			100	170	305				
KN660C	MAIZEBRAN						250	180	180	210
969502	KLC 10					100	100	100	100	100
KN718C	SUNFLOWER CAKE						90	90	95	75
KN691C	FF TOASTED SOYBEANS					30	60	80	70	70
KN778-04C	LIMESTONES 33%CA					15	50	80	95	105
	Total			1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001	Crude Protein	%	21.4	20.9	19.7	14.0	14.6	14.9	14.5	14.2
002	Crude Fat	%	4.2	3.9	3.9	4.4	7.0	6.7	6.6	6.6
060	Starch Ewers	%	35.1	38.2	38.0	40.9	39.2	37.7	37.1	37.0
003	Crude Fibre	%	6.0	3.9	4.3	5.4	6.2	6.1	6.1	5.6
089	Dry matter	%	89.3	87.5	87.5	87.8	88.4	88.9	89.1	89.1
005	Moisture	%	10.7	12.5	12.5	12.2	11.7	11.1	10.9	11.0
098	ME Poultry	Kcal	2830	2870	2820	2750	2730	2700	2640	2610
1750	Digestible Lysine	%	1.13	1.16	1.05	0.64	0.66	0.69	0.67	0.66
1768	Digestible Methion + Cystin	%	0.81	0.79	0.73	0.62	0.66	0.66	0.65	0.64
1753	Digestible Treonin	%	0.73	0.67	0.62	0.44	0.47	0.48	0.47	0.46
016	Total lysin	%	1.24	1.28	1.16	0.71	0.73	0.76	0.74	0.73
017	Total Methionin	%	0.52	0.52	0.48	0.41	0.45	0.44	0.44	0.43
019	Total Methionin + Cystin	%	0.90	0.87	0.81	0.67	0.71	0.71	0.70	0.69
048	Calcium	%	0.87	0.92	0.80	0.75	1.91	2.91	3.40	3.73
049	Total Phosphor	%	0.54	0.54	0.56	0.57	0.41	0.41	0.41	0.39
145	Digestible Phosphor	%	0.42	0.42	0.39	0.36	0.32	0.32	0.32	0.31
052	Sodium	%	0.20	0.16	0.14	0.15	0.15	0.15	0.15	0.15
050	Chloride	%	0.30	0.29	0.26	0.23	0.23	0.23	0.23	0.23
208	Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209	Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- When oil is available; replace 0,5% corn for 0,5% oil.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.

V10_08