

Mixing Instructions

With **KBC 10 SA**
 Country **Uganda**
 Specie **Broilers**
 Batch weight **1,000 kg**

9012 Optimal



Version: 5/26/2023

Code	Ingredient	Unit (kg)	969574 Galdus 0-7 days	301-9012 Starter 8-14 days	302-9012 Grower 15-28 days	303-9012 Finisher 29-42 days
			Min. 175 gr/DOC			
	969574 Galdus					
	C02627 MAIZE			440	480	580
	KN691C FF TOASTED SOYBEANS			340	340	260
	960033 KBGC 10 SA			120	100	100
	C02722 SUNFLOWER CAKE			50	50	60
	C02673 WHEAT BRAN / POLLARD			50	30	
	Total			1,000	1,000	1,000

CALCULATED ANALYSIS

001	Crude Protein	%	23.0	22.5	21.9	19.4
002	Crude Fat	%	5.4	9.0	9.0	8.1
003	Crude Fibre	%	4.0	6.2	6.0	5.7
089	Dry matter	%	89.1	89.6	89.5	89.0
004	Ash	%	5.8	5.9	5.4	5.0
098	ME Poultry	Kcal	3000	3070	3120	3150
1750	Digestible Lysine	%	1.27	1.13	1.08	0.93
1768	Digestible Methion + Cystin	%	0.86	0.86	0.81	0.76
1753	Digestible Treonin	%	0.79	0.70	0.68	0.61
016	Total lysin	%	1.40	1.28	1.23	1.06
019	Total Methionin + Cystin	%	0.95	0.96	0.91	0.85
020	Total Treonin	%	0.92	0.85	0.83	0.74
048	Calcium	%	0.86	0.82	0.70	0.69
049	Total Phosphor	%	0.54	0.52	0.49	0.46
145	Digestible Phosphor	%	0.45	0.39	0.35	0.33
052	Sodium	%	0.20	0.18	0.15	0.15
050	Chloride	%	0.29	0.31	0.27	0.27
	ZC Coccidiostat		Not Added	Added	Added	Added
	999 Essential vitamins & trace minerals		Added	Added	Added	Added

*DOC: Day Old Chick

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- When oil is available; replace 0,5% corn for 0,5% oil.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.

V10_07