

Mixing Instructions

With KBC 35 SA & KLC 20
Country Uganda
Specie Layers
Batch weight 1,000 kg

9200 -9220
Performance



Version: 5/31/2023		960710 Pre-Starter	201-9200 Starter	202-9200 Grower	204-9220 Developer	205-9220 Pre layer 18-19 weeks (till 5% production)	206-9220 Pre Peak 20-28 weeks	207-9220 Phase 1 29-55 weeks	208-9220 Phase 2 56 weeks - end
Code	Ingredient	Unit (kg)	0-7 days	1-6 weeks	7-12 weeks	13-17 weeks			
960710	Galdus Layer	Min. 100gr/DOC							
KN627C	MAIZE			525	515	555	590	600	595
969452	KBC 35/30 SA			350	300				
969564	KLC 20 + Caro					180	200	200	200
KN673C	WHEAT BRAN / POLLARD			120	180	250	150	100	85
KN778-04C	LIMESTONES 33%CA					15	55	90	105
KN768	SOYBEAN OIL			5	5		5	10	10
Total				1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001	Crude Protein	%	21.4	21.0	19.7	16.0	15.7	15.0	14.8	14.7
002	Crude Fat	%	4.2	4.3	4.4	4.0	4.3	4.6	4.6	4.1
060	Starch Ewers	%	35.1	37.0	37.3	40.5	40.9	40.6	40.3	40.0
003	Crude Fibre	%	6.0	4.0	4.3	4.7	3.9	3.5	3.3	3.3
089	Dry matter	%	89.3	87.6	87.6	87.7	88.1	88.6	88.7	88.8
005	Moisture	%	10.7	12.4	12.4	12.3	11.9	11.4	11.3	11.2
098	ME Poultry	Kcal	2830	2870	2830	2770	2790	2770	2750	2690
1750	Digestible Lysine	%	1.13	1.17	1.05	0.70	0.71	0.69	0.68	0.68
1768	Digestible Methion + Cystin	%	0.81	0.79	0.73	0.62	0.63	0.61	0.61	0.61
1753	Digestible Treonin	%	0.73	0.67	0.62	0.49	0.49	0.48	0.47	0.47
016	Total lysin	%	1.24	1.29	1.16	0.78	0.78	0.76	0.75	0.75
017	Total Methionin	%	0.52	0.52	0.48	0.40	0.41	0.40	0.40	0.40
019	Total Methionin + Cystin	%	0.90	0.87	0.81	0.69	0.69	0.67	0.66	0.66
048	Calcium	%	0.87	0.93	0.80	0.77	2.11	3.26	3.76	4.09
049	Total Phosphor	%	0.54	0.55	0.56	0.58	0.51	0.47	0.46	0.45
145	Digestible Phosphor	%	0.42	0.42	0.39	0.33	0.34	0.32	0.32	0.32
052	Sodium	%	0.20	0.16	0.14	0.14	0.15	0.15	0.15	0.15
050	Chloride	%	0.30	0.29	0.25	0.26	0.27	0.27	0.27	0.27
208	Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209	Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

- Important remarks:
- Mixing directions can change depending on the price and quality of local available raw materials.
 - Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
 - In case of no oil; replace the oil with corn. Note that FCR is likely to increase.
 - To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
 - Always make sure all birds have access to enough feed.
 - Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
 - To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
 - Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
 - In case of management related issues, ask for the Koudijs Management Guide.
 - Make sure birds always have access to fresh, cool and clean water.