

Mixing Instructions

With KBC 35/30 SA & KLC 5
Country Uganda
Specie Layers
Batch weight 1,000 kg

9200-9230
Performance



Version: 6/8/2023	Code	Ingredient	Unit (kg)	960710	201-9200	202-9200	204-9230	205-9230	206-9230	207-9230	208-9230
				Pre-Starter	Starter	Grower	Developer	Pre layer	Pre Peak	Phase 1	Phase 2
				0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	29-55 weeks	56 weeks - end
	960710	Galdus Layer	Min. 100gr/DOC								
	KN627C	MAIZE			525	515	550	580	575	585	590
	969452	KBC 35/30 SA			350	300					
	KN697C1	SOYA CAKE DEOILED					110	140	180	185	175
	KN673C	WHEAT BRAN / POLLARD			120	180	275	170	100	70	70
	KN778-04C	LIMESTONES 33%CA					15	55	85	100	110
	969501	KLC 5 + CARO					50	50	50	50	50
	KN768	SOYBEAN OIL			5	5		5	10	10	5
	Total				1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001	Crude Protein	%	21.4	21.0	19.7	15.0	15.0	15.7	15.6	15.2
002	Crude Fat	%	4.2	4.3	4.4	4.4	4.9	5.4	5.3	4.8
060	Starch Ewers	%	35.1	37.0	37.3	40.6	40.6	39.2	39.3	39.6
003	Crude Fibre	%	6.0	4.0	4.3	5.1	4.4	3.9	3.7	3.6
089	Dry matter	%	89.3	87.6	87.6	87.9	88.4	88.9	89.1	89.1
005	Moisture	%	10.7	12.4	12.4	12.1	11.6	11.1	10.9	10.9
098	ME Poultry	Kcal	2830	2870	2830	2780	2800	2800	2790	2730
1750	Digestible Lysine	%	1.13	1.17	1.05	0.64	0.67	0.74	0.74	0.71
1768	Digestible Methion + Cystin	%	0.81	0.79	0.73	0.62	0.62	0.63	0.63	0.62
1753	Digestible Treonin	%	0.73	0.67	0.62	0.46	0.47	0.50	0.50	0.49
016	Total lysin	%	1.24	1.29	1.16	0.71	0.74	0.81	0.81	0.78
017	Total Methionin	%	0.52	0.52	0.48	0.40	0.40	0.41	0.40	0.40
019	Total Methionin + Cystin	%	0.90	0.87	0.81	0.68	0.67	0.68	0.68	0.67
048	Calcium	%	0.87	0.93	0.80	0.85	2.17	3.17	3.66	3.99
049	Total Phosphor	%	0.54	0.55	0.56	0.55	0.48	0.43	0.41	0.41
145	Digestible Phosphor	%	0.42	0.42	0.39	0.35	0.33	0.32	0.32	0.31
052	Sodium	%	0.20	0.16	0.14	0.15	0.15	0.15	0.15	0.15
050	Chloride	%	0.30	0.29	0.25	0.27	0.27	0.27	0.27	0.27
208	Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209	Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC	Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

- Important remarks:
- Mixing directions can change depending on the price and quality of local available raw materials.
 - Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
 - In case of no oil; replace the oil with corn. Note that FCR is likely to increase.
 - To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
 - Always make sure all birds have access to enough feed.
 - Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
 - To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
 - Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
 - In case of management related issues, ask for the Koudijs Management Guide.
 - Make sure birds always have access to fresh, cool and clean water.