

Mixing Instructions

With KBC 35/30 SA & KLC 10
Country Uganda
Specie Layers
Batch weight 1,000 kg

9201 -9215
Optimal



Version: 6/8/2023	Code Ingredient	Unit (kg)	960710	201-9201	202-9201	204-9215	205-9215	206-9215	207-9215	208-9215
			Pre-Starter	Starter	Grower	Developer	Pre layer	Pre Peak	Phase 1	Phase 2
			0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	29-55 weeks	56 weeks - end
	960710 Galdus Layer	Min. 100gr/DOC								
	969452 KBC 35/30 SA			350	300					
	KN627C MAIZE			550	530	580	585	610	610	600
	KN673C WHEAT BRAN / POLLARD			100	170	270	190	110	95	95
	969502 KLC 10					100	100	100	100	100
	KN697C1 SOYA CAKE DEOILED					35	70	90	95	95
	KN778-04C LIMESTONES 33%CA					15	55	90	100	110
	Total			1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001 Crude Protein	%	21.4	20.9	19.7	14.1	14.5	14.4	14.4	14.4
002 Crude Fat	%	4.2	3.9	3.9	4.1	4.1	4.0	4.0	4.0
060 Starch Ewers	%	35.1	38.2	38.0	42.2	41.2	41.3	41.1	40.5
003 Crude Fibre	%	6.0	3.9	4.3	5.1	4.6	4.0	3.9	3.9
089 Dry matter	%	89.3	87.5	87.5	87.7	88.2	88.6	88.8	88.9
005 Moisture	%	10.7	12.5	12.5	12.3	11.8	11.4	11.2	11.1
098 ME Poultry	Kcal	2830	2870	2820	2770	2730	2710	2700	2660
1750 Digestible Lysine	%	1.13	1.16	1.05	0.66	0.71	0.73	0.73	0.73
1768 Digestible Methion + Cystin	%	0.81	0.79	0.73	0.63	0.63	0.63	0.63	0.63
1753 Digestible Treonin	%	0.73	0.67	0.62	0.44	0.47	0.48	0.48	0.48
016 Total lysin	%	1.24	1.28	1.16	0.72	0.77	0.79	0.79	0.79
017 Total Methionin	%	0.52	0.52	0.48	0.41	0.42	0.42	0.42	0.41
019 Total Methionin + Cystin	%	0.90	0.87	0.81	0.68	0.68	0.68	0.67	0.67
048 Calcium	%	0.87	0.92	0.80	0.75	2.07	3.23	3.56	3.89
049 Total Phosphor	%	0.54	0.54	0.56	0.56	0.50	0.45	0.44	0.43
145 Digestible Phosphor	%	0.42	0.42	0.39	0.35	0.34	0.33	0.33	0.33
052 Sodium	%	0.20	0.16	0.14	0.15	0.15	0.15	0.15	0.15
050 Chloride	%	0.30	0.29	0.26	0.23	0.23	0.23	0.23	0.23
208 Colorant added - Yellow		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
209 Colorant added - Red		Not Added	Not Added	Not Added	Added	Added	Added	Added	Added
ZC Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added

- Important remarks:
- Mixing directions can change depending on the price and quality of local available raw materials.
 - Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
 - When oil is available; replace 0,5% corn for 0,5% oil.
 - To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
 - Always make sure all birds have access to enough feed.
 - Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
 - To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
 - Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
 - In case of management related issues, ask for the Koudijs Management Guide.
 - Make sure birds always have access to fresh, cool and clean water.