

Mixing Instructions

With KBC 35/30 SA & KLC 5
Country Uganda
Specie Layers
Batch weight 1,000 kg



9201-9231
Optimal

Version: 6/8/2023		960710 Pre-Starter	201-9201 Starter	202-9201 Grower	204-9231 Developer	205-9231 Pre layer	206-9231 Pre Peak	207-9231 Phase 1	208-9231 Phase 2
		0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-28 weeks	29-55 weeks	56 weeks - end
Code	Ingredient	Unit (kg)							
960710	Galdus Layer	Min. 100gr/DOC							
KN627C	MAIZE		550	530	485	380	430	450	420
969452	KBC 35/30 SA		350	300					
KN673C	WHEAT BRAN / POLLARD		100	170	290				
KN691C	FF TOASTED SOYBEANS				110	180	190	200	190
KN660C	MAIZEBRAN					285	160	125	160
KN778-04C	LIMESTONES 33%CA				15	55	90	100	110
KN718C	SUNFLOWER CAKE				50	50	80	75	70
969501	KLC 5 + CARO				50	50	50	50	50
Total			1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001	Crude Protein	%	21.4	20.9	19.7	15.4	15.8	16.1	16.2	15.8
002	Crude Fat	%	4.2	3.9	3.9	5.9	8.6	8.0	7.8	7.8
060	Starch Ewers	%	35.1	38.2	38.0	36.7	36.4	34.7	34.5	34.0
003	Crude Fibre	%	6.0	3.9	4.3	6.5	5.3	5.8	5.6	5.5
089	Dry matter	%	89.3	87.5	87.5	88.5	88.7	89.6	89.8	89.8
005	Moisture	%	10.7	12.5	12.5	11.5	11.3	10.4	10.3	10.3
098	ME Poultry	Kcal	2830	2870	2820	2780	2800	2760	2770	2710
1750	Digestible Lysine	%	1.13	1.16	1.05	0.62	0.67	0.69	0.71	0.69
1768	Digestible Methion + Cystin	%	0.81	0.79	0.73	0.63	0.64	0.65	0.64	0.64
1753	Digestible Treonin	%	0.73	0.67	0.62	0.45	0.49	0.50	0.50	0.49
016	Total lysin	%	1.24	1.28	1.16	0.71	0.77	0.79	0.80	0.78
017	Total Methionin	%	0.52	0.52	0.48	0.41	0.43	0.43	0.43	0.42
019	Total Methionin + Cystin	%	0.90	0.87	0.81	0.69	0.71	0.71	0.71	0.70
048	Calcium	%	0.87	0.92	0.80	0.86	2.17	3.34	3.68	4.00
049	Total Phosphor	%	0.54	0.54	0.56	0.59	0.36	0.39	0.39	0.38
145	Digestible Phosphor	%	0.42	0.42	0.39	0.35	0.30	0.31	0.31	0.30
052	Sodium	%	0.20	0.16	0.14	0.15	0.16	0.16	0.16	0.16
050	Chloride	%	0.30	0.29	0.26	0.28	0.27	0.27	0.27	0.27
208	Colorant added - Yellow	Not Added	Not Added	Not Added	Added	Added	Added	Added	Added	Added
209	Colorant added - Red	Not Added	Not Added	Not Added	Added	Added	Added	Added	Added	Added
ZC	Coccidiostat	Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace minerals	Added	Added	Added	Added	Added	Added	Added	Added	Added

- Important remarks:
- Mixing directions can change depending on the price and quality of local available raw materials.
 - Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
 - When oil is available; replace 0,5% corn for 0,5% oil.
 - To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
 - Always make sure all birds have access to enough feed.
 - Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
 - To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
 - Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
 - In case of management related issues, ask for the Koudijs Management Guide.
 - Make sure birds always have access to fresh, cool and clean water.