

Mixing Instructions

With **KBC 35/30 SA & KLC 10**
 Country **Uganda** **9200-9210 Performance**
 Specie **Layers**
 Batch weight **1,000 kg**



Code Ingredient	Unit (kg)	960710	201-9200	202-9200	204-9210	205-9210	206-9210	207-9210	208-9210	209-9210
		Pre-Starter	Starter	Grower	Developer	Pre layer	Pre Peak	Phase 1	Phase 2	Phase 3
		0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-30 weeks	31-40 weeks	41-60 weeks	61 weeks - end
Version: 7/4/2023										
960710 Galdus Layer	Min. 100 g/DOC									
KN627C MAIZE			525	515	580	580	600	600	595	595
969452 KBC 35/30 SA			350	300						
KN673C WHEAT BRAN / POLLARD			120	180	270	190	100	95	95	90
969502 KLC 10					100	100	100	100	100	100
KN697C1 SOYA CAKE DEOILED					35	70	90	95	95	90
KN778-04C LIMESTONES 33%CA					15	55	100	100	110	120
KN768 SOYBEAN OIL			5	5		5	10	10	5	5
Total			1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001 Crude Protein	%	21.4	21.1	19.8	14.1	14.5	14.2	14.4	14.3	14.0
002 Crude Fat	%	4.2	4.3	4.4	4.1	4.6	5.0	5.0	4.5	4.4
003 Crude Fibre	%	6.0	3.8	4.2	5.1	4.6	3.9	3.9	3.9	3.8
089 Dry matter	%	89.3	87.7	87.7	87.7	88.3	88.9	88.9	89.0	89.1
004 Ash	%	5.9	5.9	5.5	5.1	8.9	13.1	13.1	14.1	15.0
600k ME Poultry	Kcal	2720	2790	2750	2680	2670	2670	2680	2620	2600
600 ME Poultry	MJ	11.36	11.69	11.50	11.22	11.19	11.19	11.20	10.97	10.87
1750 Digestible Lysine	%	1.13	1.18	1.06	0.65	0.70	0.71	0.72	0.72	0.71
1768 Digestible Methion + Cystin	%	0.81	0.79	0.73	0.63	0.63	0.62	0.63	0.62	0.62
048 Calcium	%	0.87	0.90	0.78	0.75	2.07	3.56	3.56	3.89	4.22
049 Total Phosphor	%	0.54	0.54	0.56	0.56	0.50	0.44	0.43	0.43	0.42
145 Digestible Phosphor	%	0.42	0.42	0.39	0.35	0.34	0.33	0.33	0.32	0.32
147 Available Phosphor	%	0.43	0.42	0.38	0.33	0.33	0.32	0.32	0.32	0.32
052 Sodium	%	0.20	0.16	0.14	0.15	0.15	0.15	0.15	0.15	0.15
Som 15 Phytase		Added	Added	Added	Added	Added	Added	Added	Added	Added
Som 16 NSP Enzyme		Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- In case of no oil; replace the oil with corn. Note that FCR is likely to increase.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- Performance mixings are focused on creating complete feed to obtain the highest performance.

V10_09