

Mixing Instructions

With **KBC 35 SA & KLC 20**
 Country **Uganda**
 Specie **Layers**
 Batch weight **1,000 kg**

9201-92211 Economy



Version: 7/4/2023

Code Ingredient	Unit (kg)	960710	201-9201	202-9201	204-92012	205-92211	206-92211	207-92211	208-92211	209-92211
		Pre-Starter	Starter	Grower	Developer	Pre layer	Pre Peak	Phase 1	Phase 2	Phase 3
		0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-30 weeks	31-40 weeks	41-60 weeks	61 weeks - end
960710 Galdus Layer	Min. 100 g/DOC									
KN627C MAIZE			550	530	555	380	450	390	390	390
969452 KBC 35/30 SA			350	300						
KN673C WHEAT BRAN / POLLARD			100	170	250					
KN660C MAIZEBRAN						350	250	310	300	290
969564 KLC 20 + Caro						180	200	200	200	200
KN778-04C LIMESTONES 33%CA					15	70	100	100	110	120
Total			1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001 Crude Protein	%	21.4	21.0	19.8	16.0	15.3	14.9	15.0	14.9	14.8
002 Crude Fat	%	4.2	3.9	3.9	4.0	6.0	5.3	5.6	5.5	5.4
003 Crude Fibre	%	6.0	3.7	4.1	4.7	3.6	3.3	3.5	3.4	3.4
089 Dry matter	%	89.3	87.6	87.6	87.6	87.6	88.1	88.0	88.2	88.3
004 Ash	%	5.9	5.9	5.5	5.3	10.8	13.5	13.6	14.6	15.5
600k ME Poultry	Kcal	2720	2800	2740	2670	2720	2660	2640	2610	2580
600 ME Poultry	MJ	11.36	11.70	11.45	11.18	11.39	11.12	11.04	10.92	10.80
1750 Digestible Lysine	%	1.13	1.18	1.06	0.70	0.68	0.67	0.67	0.67	0.66
1768 Digestible Methion + Cystin	%	0.81	0.79	0.73	0.62	0.63	0.62	0.62	0.62	0.61
048 Calcium	%	0.87	0.90	0.78	0.75	2.57	3.57	3.57	3.90	4.23
049 Total Phosphor	%	0.54	0.53	0.55	0.57	0.36	0.37	0.36	0.36	0.36
145 Digestible Phosphor	%	0.42	0.42	0.39	0.33	0.30	0.30	0.30	0.30	0.30
147 Available Phosphor	%	0.43	0.42	0.38	0.31	0.31	0.31	0.30	0.30	0.30
052 Sodium	%	0.20	0.16	0.14	0.13	0.15	0.15	0.15	0.15	0.15
Som 15 Phytase		Added								
Som 16 NSP Enzyme		Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added								

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- Economy mixings are focused on a low price per kg of complete feed and an economical performance.

V10_09