

## Mixing Instructions

With **KBC 35/30 SA & KLC 5**  
 Country **Uganda** **9200-9230 Performance**  
 Specie **Layers**  
 Batch weight **1,000 kg**



Version: 7/4/2023	960710 Pre-Starter	201-9200 Starter	202-9200 Grower	204-9230 Developer	205-9230 Pre layer 18-19 weeks (till 5% production)	206-9230 Pre Peak	207-9230 Phase 1	208-9230 Phase 2	209-9230 Phase 3
Code Ingredient	Unit (kg)	0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	20-30 weeks	31-40 weeks	41-60 weeks	61 weeks - end
960710 Galdus Layer	Min. 100 g/DOC								
KN627C MAIZE			525	515	550	580	570	590	580
969452 KBC 35/30 SA			350	300					
KN673C WHEAT BRAN / POLLARD			120	180	270	160	80	70	70
KN697C1 SOYA CAKE DEOILED					110	140	190	180	175
KN778-04C LIMESTONES 33%CA					20	65	100	100	110
969501 KLC 5 + CARO					50	50	50	50	50
KN768 SOYBEAN OIL			5	5		5	10	10	5
<b>Total</b>			<b>1,000</b>	<b>1,000</b>	<b>1,000</b>	<b>1,000</b>	<b>1,000</b>	<b>1,000</b>	<b>1,000</b>

## CALCULATED ANALYSIS

001 Crude Protein	%	21.4	21.1	19.8	14.9	14.9	15.8	15.4	15.2	15.1
002 Crude Fat	%	4.2	4.3	4.4	4.4	4.8	5.3	5.3	4.8	4.8
003 Crude Fibre	%	6.0	3.8	4.2	5.0	4.3	3.8	3.7	3.6	3.6
089 Dry matter	%	89.3	87.7	87.7	88.0	88.6	89.1	89.1	89.1	89.3
004 Ash	%	5.9	5.9	5.5	5.9	10.1	13.5	13.4	14.4	15.3
600k ME Poultry	Kcal	2720	2790	2750	2680	2700	2700	2720	2660	2630
600 ME Poultry	MJ	11.36	11.69	11.50	11.20	11.29	11.30	11.38	11.15	11.02
1750 Digestible Lysine	%	1.13	1.18	1.06	0.64	0.67	0.75	0.72	0.71	0.71
1768 Digestible Methion + Cystin	%	0.81	0.79	0.73	0.62	0.61	0.63	0.62	0.62	0.62
048 Calcium	%	0.87	0.90	0.78	1.01	2.50	3.67	3.66	3.99	4.32
049 Total Phosphor	%	0.54	0.54	0.56	0.54	0.47	0.42	0.41	0.41	0.40
145 Digestible Phosphor	%	0.42	0.42	0.39	0.34	0.33	0.32	0.31	0.31	0.31
147 Available Phosphor	%	0.43	0.42	0.38	0.32	0.31	0.31	0.31	0.31	0.31
052 Sodium	%	0.20	0.16	0.14	0.15	0.15	0.15	0.15	0.15	0.15
Som 15 Phytase		Added	Added	Added	Added	Added	Added	Added	Added	Added
Som 16 NSP Enzyme		Added	Not Added	Not Added	Added	Added	Added	Added	Added	Added
ZC Coccidiostat		Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999 Essential vitamins & trace minerals		Added	Added	Added	Added	Added	Added	Added	Added	Added

### Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- In case of no oil; replace the oil with corn. Note that FCR is likely to increase.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- Performance mixings are focused on creating complete feed to obtain the highest performance.

V10\_09