

Mixing Instructions

With **KBC 35 SA & KLC 10**

Country **Uganda**

Specie **Layers**

Batch weight **1,000 kg**

9201-92151 Economy



Version: 10/23/2023

Code	Ingredient	Unit (kg)	960710	201-9201	202-9201	204-92011	205-92151	206-92151	207-92151	208-92151	209-92151
			Pre-Starter	Starter	Grower	Developer	Pre layer	Pre Peak	Phase 1	Phase 2	Phase 3
			0-7 days	1-6 weeks	7-12 weeks	13-17 weeks	18-19 weeks (till 5% production)	20-30 weeks	31-40 weeks	41-60 weeks	61 weeks - end
960710	Galdus Layer	Min. 100 g/DOC									
KN627C	MAIZE			550	530	580	520	550	460	460	460
KN649C	MAIZEBRAN						100	100	200	210	210
969452	KBC 35/30 SA			350	300						
KN673C	WHEAT BRAN / POLLARD			100	170	270	150	50	50	30	20
969502	KLC 10					100	100	100	100	100	100
KN697C1	SOYA CAKE DEOILED					35	70	100	90	90	90
KN778-04C	LIMESTONES 33%CA					15	60	100	100	110	120
Total				1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000

CALCULATED ANALYSIS

001	Crude Protein	%	21.1	20.9	19.7	14.1	14.4	14.5	14.4	14.2	14.0
002	Crude Fat	%	4.0	3.9	3.9	4.2	4.8	4.7	5.3	5.3	5.3
003	Crude Fibre	%	5.8	3.9	4.3	5.1	4.5	3.8	4.0	3.8	3.8
089	Dry matter	%	89.1	87.6	87.6	87.7	88.2	88.6	88.5	88.6	88.7
004	Ash	%	5.8	5.8	5.5	5.1	9.5	13.2	13.4	14.3	15.3
600k	ME Poultry	Kcal	2760	2790	2730	2690	2660	2630	2630	2630	2610
600	ME Poultry	MJ	11.53	11.68	11.43	11.24	11.14	11.14	11.02	11.00	10.92
1750	Digestible Lysine	%	1.13	1.16	1.05	0.65	0.70	0.74	0.72	0.71	0.71
1768	Digestible Methion + Cystin	%	0.81	0.79	0.73	0.63	0.64	0.64	0.64	0.63	0.63
048	Calcium	%	0.87	0.90	0.78	0.75	2.23	3.56	3.56	3.88	4.21
049	Total Phosphor	%	0.53	0.54	0.56	0.56	0.46	0.40	0.38	0.36	0.35
145	Digestible Phosphor	%	0.42	0.42	0.39	0.35	0.33	0.32	0.31	0.31	0.31
147	Available Phosphor	%	0.44	0.42	0.38	0.34	0.33	0.32	0.31	0.31	0.31
052	Sodium	%	0.20	0.16	0.14	0.15	0.15	0.15	0.15	0.15	0.15
Som 15	Phytase	Added	Added	Added	Added	Added	Added	Added	Added	Added	Added
Som 16	NSP Enzyme	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
ZC	Coccidiostat	Not Added	Added	Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added	Not Added
999	Essential vitamins & trace	Added	Added	Added	Added	Added	Added	Added	Added	Added	Added

Important remarks:

- Mixing directions can change depending on the price and quality of local available raw materials.
- Consult the local legislation for possible regulations regarding use of feed additives such as coccidiostats and withdrawal time.
- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- In case of management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.
- Optimal mixings are focused on revenue. With this mixing you create complete feed for optimal performance and optimal revenue.

V10_09