

## Mixing Instructions

Client **Uganda**  
 With **KLC 20**  
 Intended for **Indigenous - Laying prod.**  
 Batch weight **1.000 kg**



| Version: | 27-8-2019            |           | EAfrica.201<br>Starter | EAfrica.202<br>Grower | EAfrica.203<br>Developer | EAfrica.204<br>Prelay | EAfrica.205<br>Layer | EAfrica.206<br>Layer 2 |
|----------|----------------------|-----------|------------------------|-----------------------|--------------------------|-----------------------|----------------------|------------------------|
| Code     | Ingredient           | Unit (kg) | 0-6 wks                | 7-12 wks              | 13-17 wks                | 17 wks-5% prod        | 5% prod-40 wks       | >40 wks                |
| 969564   | KLC 20 + Caro        |           | 250                    | 200                   | 180                      | 200                   | 200                  | 200                    |
| C02627   | MAIZE 8,5%CP         |           | 400                    | 455                   | 480                      | 535                   | 550                  | 530                    |
| C02673   | WHEAT BRAN           |           | 30                     | 35                    | 220                      | 160                   | 65                   | 70                     |
| C02778   | LIMESTONES           |           | 15                     | 15                    | 20                       | 55                    | 85                   | 100                    |
| KN660C   | MAIZEBRAN            |           | 150                    | 250                   | 100                      | 50                    | 100                  | 100                    |
| KN691C   | FF TOASTED SOYBEANS  |           | 150                    | 40                    |                          |                       |                      |                        |
| KN788    | DICALPHOS 18,5P/26CA |           | 5                      | 5                     |                          |                       |                      |                        |
|          | <b>Total</b>         |           | <b>1.000</b>           | <b>1.000</b>          | <b>1.000</b>             | <b>1.000</b>          | <b>1.000</b>         | <b>1.000</b>           |

## CALCULATED ANALYSIS

|     |                                     |       |       |       |       |       |       |       |
|-----|-------------------------------------|-------|-------|-------|-------|-------|-------|-------|
| 001 | Crude Protein                       | %     | 22,0  | 17,2  | 16,2  | 16,1  | 15,3  | 15,2  |
| 002 | Crude Fat                           | %     | 7,0   | 6,1   | 4,7   | 4,2   | 4,4   | 4,4   |
| 003 | Crude Fibre                         | %     | 4,2   | 3,8   | 4,4   | 3,9   | 3,3   | 3,3   |
| 098 | ME Poultry                          | Kcal  | 2857  | 2800  | 2704  | 2699  | 2684  | 2626  |
| 016 | Total lysin                         | %     | 1,22  | 0,88  | 0,79  | 0,81  | 0,77  | 0,77  |
| 017 | Total Methionin                     | %     | 0,53  | 0,44  | 0,40  | 0,41  | 0,41  | 0,40  |
| 019 | Total Methionin + Cystin            | %     | 0,91  | 0,75  | 0,70  | 0,71  | 0,69  | 0,68  |
| 020 | Total Treonin                       | %     | 0,83  | 0,64  | 0,59  | 0,59  | 0,57  | 0,56  |
| 048 | Calcium                             | %     | 0,93  | 0,86  | 0,88  | 2,06  | 3,06  | 3,56  |
| 049 | Total Phosphor                      | %     | 0,57  | 0,50  | 0,53  | 0,50  | 0,42  | 0,42  |
| 052 | Sodium                              | %     | 0,19  | 0,15  | 0,14  | 0,15  | 0,15  | 0,15  |
| 050 | Chloride                            | %     | 0,32  | 0,27  | 0,25  | 0,27  | 0,26  | 0,26  |
| 999 | Essential vitamins & trace minerals | IU/kg | Added | Added | Added | Added | Added | Added |

Note: Mixing directions can change depending on the price and quality of local available raw materials.

For the best results; start with the Koudijs Galdus 2mm pre-starter (minimum 150 gram/bird total, preferably more).

### Important remarks:

- To achieve good results, follow up on the mixing instructions! Do not alter or change without consult.
- Always make sure all birds have access to enough feed.
- Ensure proper quality of the limestone used. Quicktests are available, ask your local representative for more information.
- To avoid drop in feed intake; ensure the change of feeds (from Starter to Grower feed etc.) happens gradually.
- Be biosecurity wise; use disinfection materials, keep separate shoes for the bird houses only.
- For more management related issues, ask for the Koudijs Management Guide.
- Make sure birds always have access to fresh, cool and clean water.

V09\_05

